

ON APPROXIMATE CONNES-AMENABILITY OF ENVELOPING DUAL BANACH ALGEBRAS

AHMAD SHIRINKALAM AND A. POURABBAS

ABSTRACT. For a Banach algebra $\mathcal{A}$, we introduce various approximate virtual diagonals such as approximate WAP-virtual diagonal and approximate virtual diagonal.

For the enveloping dual Banach algebra $F(\mathcal{A})$ of $\mathcal{A}$, we show that $F(\mathcal{A})$ is approximately Connes-amenable if and only if $\mathcal{A}$ has an approximate WAP-virtual diagonal.

Further, for a discrete group G , we show that if the group algebra $\ell^1(G)$ has an approximate WAP-virtual diagonal, then it has an approximate virtual diagonal.

1. INTRODUCTION AND PRELIMINARIES

The notion of approximate Connes-amenability for a dual Banach algebra was first introduced by Esslamzadeh *et al.* (2012). A dual Banach algebra $\mathcal{A}$ is called approximately Connes-amenable if for every normal dual $\mathcal{A}$ -bimodule X , every w^* -continuous derivation $D : \mathcal{A} \rightarrow X$ is approximately inner, that is, there exists a net $(x_\alpha) \subseteq X$ such that $D(b) = \lim_\alpha b \cdot x_\alpha - x_\alpha \cdot b$ ($b \in \mathcal{A}$).

Let $\mathcal{A}$ be a Banach algebra and let X be a Banach $\mathcal{A}$ -bimodule. An element $x \in X$ is called weakly almost periodic if the module maps $\mathcal{A} \rightarrow X$; $a \mapsto a \cdot x$ and $a \mapsto x \cdot a$ are weakly compact. The set of all weakly almost periodic elements of X is denoted by $\text{WAP}(X)$ which is a norm-closed subspace of X .

For a Banach algebra $\mathcal{A}$, one of the most important subspaces of $\mathcal{A}^*$ is $\text{WAP}(\mathcal{A}^*)$ which is left introverted in the sense of [6, §1]. Runde (2004) observed that, the space $F(\mathcal{A}) = \text{WAP}(\mathcal{A}^*)^*$ is a dual Banach algebra with the first Arens product inherited from $\mathcal{A}^{**}$. He also showed that $F(\mathcal{A})$ is a *canonical* dual Banach algebra associated to $\mathcal{A}$ (see [6] or [8, Theorem 4.10] for more details).

Choi *et al.* (2014) called $F(\mathcal{A})$ the enveloping dual Banach algebra associated to $\mathcal{A}$ and they studied Connes-amenability of $F(\mathcal{A})$. Indeed, they introduced the notion of WAP-virtual diagonal for a Banach algebra $\mathcal{A}$ and they showed that for a given Banach algebra $\mathcal{A}$, the dual Banach algebra $F(\mathcal{A})$ is Connes-amenable if and only if $\mathcal{A}$ admits a WAP-virtual diagonal. They also showed that for a group algebra $L^1(G)$, the existence of a virtual diagonal is equivalent to the existence of a WAP-virtual diagonal. As a consequent $L^1(G)$ is amenable if and only if $F(L^1(G))$ is Connes-amenable, a fact that previously was shown by Runde (2004).

2010 *Mathematics Subject Classification.* Primary: 46H20; Secondary: 46H25, 46H35.

Key words and phrases. Enveloping dual Banach algebra, Approximately Connes-amenable, Approximate WAP-virtual diagonal, Approximate virtual diagonal.

Motivated by these results, we investigate an *approximate* analogue of WAP-virtual diagonal related to approximate Conns-amenability of $F(\mathcal{A})$. Indeed, we introduce a notion of an approximate WAP-virtual diagonal for $\mathcal{A}$ and we show that $F(\mathcal{A})$ is approximately Connes-amenable if and only if $\mathcal{A}$ has an approximate WAP-virtual diagonal. We also introduce various notions of approximate-type virtual diagonals for a Banach algebra -such as approximate virtual diagonal- and we show that for a discrete group G , if $\ell^1(G)$ has an approximate WAP-virtual diagonal, then it has an approximate virtual diagonal.

Given a Banach algebra $\mathcal{A}$, its unitization is denoted by $\mathcal{A}^\#$. For a Banach $\mathcal{A}$ -bimodule X , the topological dual space X^* of X becomes a Banach $\mathcal{A}$ -bimodule via the following actions

$$(1.1) \quad \langle x, a \cdot \varphi \rangle = \langle x \cdot a, \varphi \rangle, \quad \langle x, \varphi \cdot a \rangle = \langle a \cdot x, \varphi \rangle \quad (a \in \mathcal{A}, x \in X, \varphi \in X^*).$$

Definition 1.1. A Banach algebra $\mathcal{A}$ is called dual, if it is a dual Banach space with a predual $\mathcal{A}_*$ such that the multiplication in $\mathcal{A}$ is separately $\sigma(\mathcal{A}, \mathcal{A}_*)$ -continuous [7, Definition 1.1]. Equivalently, a Banach algebra $\mathcal{A}$ is dual if it has a (not necessarily unique) predual $\mathcal{A}_*$ which is a closed submodule of $\mathcal{A}^*$ [9, Exercise 4.4.1].

Let $\mathcal{A}$ be a dual Banach algebra and let a dual Banach space X be an $\mathcal{A}$ -bimodule. An element $x \in X$ is called normal, if the module actions $a \mapsto a \cdot x$ and $a \mapsto x \cdot a$ are w^* - w^* -continuous. The set of all normal elements in X is denoted by X_σ . We say that X is normal if $X = X_\sigma$.

Recall that for a Banach algebra $\mathcal{A}$ and a Banach $\mathcal{A}$ -bimodule X , a bounded linear map $D : \mathcal{A} \rightarrow X$ is called a bounded derivation if $D(ab) = a \cdot D(b) + D(a) \cdot b$ for every $a, b \in \mathcal{A}$. A derivation $D : \mathcal{A} \rightarrow X$ is called inner if there exists an element $x \in X$ such that for every $a \in \mathcal{A}$, $D(a) = \text{ad}_x(a) = a \cdot x - x \cdot a$. A dual Banach algebra $\mathcal{A}$ is called Connes-amenable if for every normal dual $\mathcal{A}$ -bimodule X , every w^* - w^* -continuous derivation $D : \mathcal{A} \rightarrow X$ is inner [8, Definition 1.5].

Following [2], for a given Banach algebra $\mathcal{A}$ and a Banach $\mathcal{A}$ -bimodule X , the $\mathcal{A}$ -bimodule $\text{WAP}(X^*)$ is denoted by $F_{\mathcal{A}}(X)_*$ and its dual space $\text{WAP}(X^*)^*$ is denoted by $F_{\mathcal{A}}(X)$. Choi *et al.* [2, Theorem 4.3] showed that if X is a Banach $\mathcal{A}$ -bimodule, then $F_{\mathcal{A}}(X)$ is a normal dual $F(\mathcal{A})$ -bimodule. Since $\text{WAP}(X^*)$ is a closed subspace of X^* , we have a quotient map $q : X^{**} \twoheadrightarrow F_{\mathcal{A}}(X)$. Composing the canonical inclusion map $X \hookrightarrow X^{**}$ with q , we obtain a continuous $\mathcal{A}$ -bimodule map $\eta_X : X \rightarrow F_{\mathcal{A}}(X)$ which has a w^* -dense range. In the special case where $X = \mathcal{A}$, we usually omit the subscript and simply use the notation $F(\mathcal{A})$ and the map $\eta_{\mathcal{A}} : \mathcal{A} \rightarrow F(\mathcal{A})$ is an algebra homomorphism.

It is well-known that for a given Banach algebra $\mathcal{A}$, the projective tensor product $\mathcal{A} \hat{\otimes} \mathcal{A}$ is a Banach $\mathcal{A}$ -bimodule through

$$a \cdot (b \otimes c) := ab \otimes c, \quad (b \otimes c) \cdot a := b \otimes ca, \quad (a, b, c \in \mathcal{A})$$

and the map $\Delta : \mathcal{A} \hat{\otimes} \mathcal{A} \rightarrow \mathcal{A}$ defined on elementary tensors by $\Delta(a \otimes b) = ab$ and extended by linearity and continuity, is an $\mathcal{A}$ -bimodule map with respect to the module structure of $\mathcal{A} \hat{\otimes} \mathcal{A}$. Let $\Delta_{\text{WAP}} : F_{\mathcal{A}}(\mathcal{A} \hat{\otimes} \mathcal{A}) \rightarrow F(\mathcal{A})$ be the w^* - w^* -continuous $\mathcal{A}$ -bimodule map induced by $\Delta : \mathcal{A} \hat{\otimes} \mathcal{A} \rightarrow \mathcal{A}$ (see [2, Corollary 5.2] for more details).

Definition 1.2. [2, Definition 6.4] An element $M \in F_{\mathcal{A}}(\mathcal{A} \hat{\otimes} \mathcal{A})$ is called a WAP-virtual diagonal for $\mathcal{A}$ if

$$a \cdot M = M \cdot a \quad \text{and} \quad \Delta_{\text{WAP}}(M) \cdot a = \eta_{\mathcal{A}}(a) \quad (a \in \mathcal{A}).$$

Recall that an $\mathcal{A}$ -bimodule X is called neo-unital if every $x \in X$ can be written as $a \cdot y \cdot b$ for some $a, b \in \mathcal{A}$ and $y \in X$.

Remark 1.3. Let $\mathcal{A}$ be a Banach algebra with a bounded approximate identity and let X be a Banach $\mathcal{A}$ -bimodule. Then by [5, Proposition 1.8], the subspace $X_{ess} := \text{lin}\{a \cdot x \cdot b : a, b \in \mathcal{A}, x \in X\}$ is a neo-unital closed sub- $\mathcal{A}$ -bimodule of X . Moreover, $X_{ess}^\perp$ is complemented in X^* .

2. APPROXIMATE CONNES-AMENABILITY OF $F(\mathcal{A})$

In this section we find some conditions under which $F(\mathcal{A})$ is approximately Connes-amenable. For an arbitrary dual Banach algebra $\mathcal{A}$, approximate Connes-amenable of $\mathcal{A}$ is equivalent to approximate Connes-amenable of $\mathcal{A}^\#$ [3, Proposition 2.3], so without loss of generality throughout of this section we may suppose that $F(\mathcal{A})$ has an identity element e .

Remark 2.1. Let $\mathcal{A}$ be a Banach algebra with an identity element e . Let X be a normal dual $\mathcal{A}$ -bimodule and let $D : \mathcal{A} \rightarrow X$ be a bounded derivation. By [4, Lemma 2.3], there exists a bounded derivation $D_1 : \mathcal{A} \rightarrow e \cdot X \cdot e$ such that $D = D_1 + \text{ad}_x$ for some $x \in X$. Furthermore, D is inner derivation if and only if D_1 is inner.

Lemma 2.2. *Let $\mathcal{A}$ be a Banach algebra. Then $F(\mathcal{A})$ is approximately Connes-amenable if and only if every bounded derivation from $\mathcal{A}$ into a unit-linked, normal dual $F(\mathcal{A})$ -bimodule is approximately inner.*

Proof. Suppose that $F(\mathcal{A})$ is approximately Connes-amenable. Let N be a unit-linked, normal dual $F(\mathcal{A})$ -bimodule and let $D : \mathcal{A} \rightarrow N$ be a bounded derivation. We show that D is approximately inner. By [3, Theorem 4.4], there is a w^* - w^* -continuous derivation $\tilde{D} : F(\mathcal{A}) \rightarrow F_{\mathcal{A}}(N)$ such that $\tilde{D}\eta_{\mathcal{A}} = \eta_N D$, that is, the following diagram commutes

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{D} & N \\ \eta_{\mathcal{A}} \downarrow & & \downarrow \eta_N \\ F(\mathcal{A}) & \xrightarrow{\tilde{D}} & F_{\mathcal{A}}(N). \end{array}$$

Since $F_{\mathcal{A}}(N)$ is a normal dual $F(\mathcal{A})$ -bimodule and $F(\mathcal{A})$ is approximately Connes-amenable, the derivation $\tilde{D}$ is approximately inner. Since $\eta_{\mathcal{A}}$ is an algebra homomorphism, the derivation $\tilde{D}\eta_{\mathcal{A}}$ is also approximately inner. Hence the derivation $\eta_N D$ is approximately inner, that is, there is a net $(\xi_i) \subseteq F_{\mathcal{A}}(N)$ such that

$$\eta_N D(a) = \lim_i a \cdot \xi_i - \xi_i \cdot a \quad (a \in \mathcal{A}).$$

By [3, Corollary 5.3] there is a w^* - w^* -continuous $\mathcal{A}$ -bimodule map $\epsilon_N : F_{\mathcal{A}}(N) \rightarrow N$ such that $\epsilon_N \eta_N = \text{id}_N$. Now by setting $x_i = \epsilon_N(\xi_i) \in N$ for each i we have

$$D(a) = \epsilon_N \eta_N D(a) = \epsilon_N (\lim_i a \cdot \xi_i - \xi_i \cdot a) = \lim_i a \cdot x_i - x_i \cdot a,$$

so $D : \mathcal{A} \rightarrow N$ is approximately inner.

Conversely, suppose that every bounded derivation from $\mathcal{A}$ into a unit-linked, normal dual $F(\mathcal{A})$ -bimodule is approximately inner. Let N be a normal dual $F(\mathcal{A})$ -bimodule and let $D : F(\mathcal{A}) \rightarrow N$ be a w^* - w^* -continuous derivation. By Remark 2.1 without loss of generality we may assume that N is a unit-linked, normal dual $F(\mathcal{A})$ -bimodule. We shall show that D is approximately inner. Since $\eta_{\mathcal{A}}$ is an algebra homomorphism, the map $D' := D\eta_{\mathcal{A}} : \mathcal{A} \rightarrow N$ is a bounded derivation, so by assumption D' is approximately inner. Since N is normal, D' extends to a w^* - w^* -continuous approximately inner derivation $D'' : F(\mathcal{A}) \rightarrow N$. The derivations D and D'' are w^* - w^* -continuous and they agree on the w^* -dense subset $\eta_{\mathcal{A}}(\mathcal{A}) \subseteq F(\mathcal{A})$, so they agree on $F(\mathcal{A})$. Hence D is approximately inner, that is, $F(\mathcal{A})$ is approximately Connes-amenable. $\square$

Remark 2.3. Since $F(\mathcal{A})$ has an identity element e , by [2, Theorem 6.8] there exists an element $s \in F_{\mathcal{A}}(\mathcal{A} \hat{\otimes} \mathcal{A})$ such that $\Delta_{\text{WAP}}(s) = e$. We use this fact in the following lemma several times.

Lemma 2.4. *Let $\mathcal{A}$ be a Banach algebra. Then the following are equivalent*

- (i) $F(\mathcal{A})$ is approximately Connes-amenable;
- (ii) *There exists a net $(M_{\alpha})_{\alpha} \subseteq F_{\mathcal{A}}(\mathcal{A} \hat{\otimes} \mathcal{A})$ such that for every $a \in \mathcal{A}$, $a \cdot M_{\alpha} - M_{\alpha} \cdot a \rightarrow 0$ and $\Delta_{\text{WAP}}(M_{\alpha}) = e$ for all α .*

Proof. (i) $\Rightarrow$ (ii) Fix $s \in F_{\mathcal{A}}(\mathcal{A} \hat{\otimes} \mathcal{A})$ as mentioned in Remark 2.3. We note that $\ker \Delta_{\text{WAP}}$ is a normal dual $F(\mathcal{A})$ -bimodule, now we define a w^* - w^* -continuous derivation $D : F(\mathcal{A}) \rightarrow \ker \Delta_{\text{WAP}}$ by $D(\mathbf{a}) = s \cdot \mathbf{a} - \mathbf{a} \cdot s$ ($\mathbf{a} \in F(\mathcal{A})$). Since $F(\mathcal{A})$ is approximately Connes-amenable, the derivation D is approximately inner. Thus there is a net $(N_{\alpha}) \subseteq \ker \Delta_{\text{WAP}}$ such that for every $a \in \mathcal{A}$

$$(2.1) \quad s \cdot a - a \cdot s = \lim_{\alpha} (a \cdot N_{\alpha} - N_{\alpha} \cdot a).$$

If we set $M_{\alpha} = N_{\alpha} + s$ for every α , then we have $\Delta_{\text{WAP}}(M_{\alpha}) = \Delta_{\text{WAP}}(s) = e$ and for every $a \in \mathcal{A}$ using (2.1) we have

$$a \cdot M_{\alpha} - M_{\alpha} \cdot a \rightarrow 0,$$

as required.

(ii) $\Rightarrow$ (i) The hypothesis in (ii) ensures that each $M_{\alpha} - s$ is in $\ker \Delta_{\text{WAP}}$ and for every $a \in \mathcal{A}$

$$(2.2) \quad a \cdot (M_{\alpha} - s) - (M_{\alpha} - s) \cdot a \rightarrow s \cdot a - a \cdot s.$$

Let N be a unit-linked normal dual $F(\mathcal{A})$ -bimodule and let $D : \mathcal{A} \rightarrow N$ be a bounded derivation. We show that D is approximately inner, so by Lemma 2.2, $F(\mathcal{A})$ is approximately Connes-amenable. Using the terminology of [2, Theorem 6.8], if we define $d(a) = s \cdot a - a \cdot s$ for all $a \in \mathcal{A}$, then $d : \mathcal{A} \rightarrow \ker \Delta_{\text{WAP}}$ is weakly universal for derivation D with coefficient in N , that is, there exists a w^* - w^* -continuous (and so it is norm continuous) $\mathcal{A}$ -bimodule map $f : \ker \Delta_{\text{WAP}} \rightarrow N$ such that $fd = D$. Set $y_{\alpha} = f(M_{\alpha} - s)$ for every α . Using (2.2) for every $a \in \mathcal{A}$ we have

$$\begin{aligned} D(a) &= fd(a) = f(s \cdot a - a \cdot s) \\ &= f(\lim_{\alpha} a \cdot (M_{\alpha} - s) - (M_{\alpha} - s) \cdot a) \\ &= \lim_{\alpha} f(a \cdot (M_{\alpha} - s) - (M_{\alpha} - s) \cdot a) \\ &= \lim_{\alpha} a \cdot y_{\alpha} - y_{\alpha} \cdot a. \end{aligned}$$

Hence D is approximately inner and this completes the proof. $\square$

Now we introduce a notion of an approximate WAP-virtual diagonal for $\mathcal{A}$.

Definition 2.5. Let $\mathcal{A}$ be a Banach algebra and let $F(\mathcal{A})$ has an identity element e . An *approximate WAP-virtual diagonal* for $\mathcal{A}$ is a net $(M_\alpha)_\alpha \subseteq F_{\mathcal{A}}(\mathcal{A} \hat{\otimes} \mathcal{A})$ such that

$$(2.3) \quad a \cdot M_\alpha - M_\alpha \cdot a \rightarrow 0 \quad (a \in \mathcal{A}) \quad \text{and} \quad \Delta_{\text{WAP}}(M_\alpha) \rightarrow e$$

in the norm topology.

Theorem 2.6. Let $\mathcal{A}$ be a Banach algebra. Then the following are equivalent

- (i) $F(\mathcal{A})$ is approximately Connes-amenable;
- (ii) $\mathcal{A}$ has an approximate WAP-virtual diagonal.

Proof. (i) $\Rightarrow$ (ii) It is immediate by Lemma 2.4.

(ii) $\Rightarrow$ (i) Suppose that $\mathcal{A}$ has an approximate WAP-virtual diagonal. Then there exists a net $(M_\alpha)_\alpha \subseteq F_{\mathcal{A}}(\mathcal{A} \hat{\otimes} \mathcal{A})$ such that for every $a \in \mathcal{A}$,

$$(2.4) \quad a \cdot M_\alpha - M_\alpha \cdot a \rightarrow 0 \quad \text{and} \quad \Delta_{\text{WAP}}(M_\alpha) \rightarrow e.$$

Let N be a unit-linked normal dual $F(\mathcal{A})$ -bimodule and let $D : \mathcal{A} \rightarrow N$ be a bounded derivation. We show that D is approximately inner and then by Lemma 2.2, $F(\mathcal{A})$ is approximately Connes-amenable.

For every $x \in N_*$ consider the functional $f_x \in F_{\mathcal{A}}(\mathcal{A} \hat{\otimes} \mathcal{A})^*$ defined by $f_x(a \otimes b) = (a \cdot D(b))(x) \quad (a, b \in \mathcal{A})$. Note that for every $m \in \mathcal{A} \hat{\otimes} \mathcal{A}$ we have

$$f_{x \cdot a - a \cdot x}(m) = (f_x \cdot a - a \cdot f_x)(m) + (\Delta(m) \cdot D(a))(x).$$

Consider the $\mathcal{A}$ -bimodule map $\Delta : \mathcal{A} \hat{\otimes} \mathcal{A} \rightarrow \mathcal{A}$. Then by [2, Corollary 5.2] there exists a unique w^* - w^* -continuous linear map $\Delta_{\text{WAP}} : F_{\mathcal{A}}(\mathcal{A} \hat{\otimes} \mathcal{A}) \rightarrow F(\mathcal{A})$ makes the following diagram

$$\begin{array}{ccc} \mathcal{A} \hat{\otimes} \mathcal{A} & \xrightarrow{\Delta} & \mathcal{A} \\ \eta_{\mathcal{A} \hat{\otimes} \mathcal{A}} \downarrow & & \downarrow \eta_{\mathcal{A}} \\ F_{\mathcal{A}}(\mathcal{A} \hat{\otimes} \mathcal{A}) & \xrightarrow{\Delta_{\text{WAP}}} & F(\mathcal{A}) \end{array}$$

commute. Since $\eta_{\mathcal{A} \hat{\otimes} \mathcal{A}}(\mathcal{A} \hat{\otimes} \mathcal{A})$ is w^* -dense in $F_{\mathcal{A}}(\mathcal{A} \hat{\otimes} \mathcal{A})$, there is a net $(m_\beta^\alpha)_\beta \subseteq \mathcal{A} \hat{\otimes} \mathcal{A}$ such that $\eta_{\mathcal{A} \hat{\otimes} \mathcal{A}}(m_\beta^\alpha) \xrightarrow{w^*} M_\alpha$ for every α .

Set $\lambda_\alpha(x) = \langle M_\alpha, f_x \rangle$ for each α , we have

$$\begin{aligned}
\langle a \cdot \lambda_\alpha - \lambda_\alpha \cdot a, x \rangle &= \langle \lambda_\alpha, x \cdot a - a \cdot x \rangle \\
&= \langle M_\alpha, f_{x \cdot a - a \cdot x} \rangle \\
&= \lim_\beta \langle \eta_{\mathcal{A} \hat{\otimes} \mathcal{A}}(m_\beta^\alpha), f_{x \cdot a - a \cdot x} \rangle \\
&= \lim_\beta \langle f_x \cdot a - a \cdot f_x, m_\beta^\alpha \rangle + \lim_\beta \langle \Delta(m_\beta^\alpha) \cdot D(a), x \rangle \\
&= \lim_\beta \langle f_x \cdot a - a \cdot f_x, m_\beta^\alpha \rangle + \lim_\beta \langle x, \eta_{\mathcal{A}} \Delta(m_\beta^\alpha) \cdot D(a) \rangle \\
&= \lim_\beta \langle \eta_{\mathcal{A} \hat{\otimes} \mathcal{A}}(m_\beta^\alpha), f_x \cdot a - a \cdot f_x \rangle + \lim_\beta \langle x, \Delta_{\text{WAP}} \eta_{\mathcal{A} \hat{\otimes} \mathcal{A}}(m_\beta^\alpha) \cdot D(a) \rangle \\
&= \langle a \cdot M_\alpha - M_\alpha \cdot a, f_x \rangle + \langle x, \Delta_{\text{WAP}}(M_\alpha) \cdot D(a) \rangle.
\end{aligned}$$

Because $\|f_x\| \leq \|D\| \|x\|$,

$$\|(a \cdot \lambda_\alpha - \lambda_\alpha \cdot a)(x) - D(a)(x)\| \leq \|a \cdot M_\alpha - M_\alpha \cdot a\| \|D\| \|x\| + \|x\| \|\Delta_{\text{WAP}}(M_\alpha) - e\| \|D(a)\|.$$

This together with equation (2.4) imply that $D(a) = \lim_\alpha \text{ad}_{\lambda_\alpha}(a)$ for every $a \in \mathcal{A}$, so D is approximately inner. $\square$

3. APPROXIMATE-TYPE VIRTUAL DIAGONALS

Suppose that $\mathcal{A}$ is a Banach algebra. Consider the $\mathcal{A}$ -bimodule map $\Delta^* : \mathcal{A}^* \rightarrow (\mathcal{A} \hat{\otimes} \mathcal{A})^*$. Let $V \subseteq (\mathcal{A} \hat{\otimes} \mathcal{A})^*$ be a closed subspace and let $E \subseteq (\Delta^*)^{-1}(V) \subseteq \mathcal{A}^*$. Then we obtain a map $\Delta^*|_E : E \rightarrow V$. We denote the adjoint of $\Delta^*|_E$ by $\Delta_E : V^* \rightarrow E^*$.

Definition 3.1. Let $V \subseteq (\mathcal{A} \hat{\otimes} \mathcal{A})^*$ be a non-zero closed subspace and let $E \subseteq (\Delta^*)^{-1}(V) \subseteq \mathcal{A}^*$. An *approximate V -virtual diagonal* for $\mathcal{A}$ is a net $(M_\alpha) \subseteq V^*$ satisfies

$$a \cdot M_\alpha - M_\alpha \cdot a \rightarrow 0 \quad \text{and} \quad \langle \Delta_E(M_\alpha) \cdot a, \varphi \rangle \rightarrow \langle \varphi, a \rangle \quad (a \in \mathcal{A}, \varphi \in E).$$

We consider the following cases

- 1- Let $V = (\mathcal{A} \hat{\otimes} \mathcal{A})^*$. Then we obtain a new concept of a diagonal for $\mathcal{A}$, called an *approximate virtual diagonal*, that is, a -not necessarily bounded- net $(M_\alpha) \subseteq (\mathcal{A} \hat{\otimes} \mathcal{A})^{**}$ such that

$$a \cdot M_\alpha - M_\alpha \cdot a \rightarrow 0 \quad \text{and} \quad \langle \Delta^{**}(M_\alpha) \cdot a, \varphi \rangle \rightarrow \langle \varphi, a \rangle \quad (a \in \mathcal{A}, \varphi \in \mathcal{A}^*).$$

- 2- Let $V = (\mathcal{A} \hat{\otimes} \mathcal{A})_{ess}^* \subseteq (\mathcal{A} \hat{\otimes} \mathcal{A})^*$. Then we obtain an *approximate $(\mathcal{A} \hat{\otimes} \mathcal{A})_{ess}^*$ -virtual diagonal* for $\mathcal{A}$.

In the following proposition we show that how these two diagonals deal with.

Proposition 3.2. *Let $\mathcal{A}$ be a Banach algebra with a bounded approximate identity. If $\mathcal{A}$ has an approximate $(\mathcal{A} \hat{\otimes} \mathcal{A})_{ess}^*$ -virtual diagonal, then it has an approximate virtual diagonal.*

Proof. Let $V = (\mathcal{A} \hat{\otimes} \mathcal{A})_{ess}^*$. By Remark 1.3, V is a closed subspace of $(\mathcal{A} \hat{\otimes} \mathcal{A})^*$ and we have an isomorphism $\tau : V^* \rightarrow (\mathcal{A} \hat{\otimes} \mathcal{A})^{**}/V^\perp$ defined by $\tau(T) = x + V^\perp$ for some $x \in (\mathcal{A} \hat{\otimes} \mathcal{A})^{**}$. Since $V^\perp$ is complemented in $(\mathcal{A} \hat{\otimes} \mathcal{A})^{**}$, there exists an $\mathcal{A}$ -bimodule projection $P : (\mathcal{A} \hat{\otimes} \mathcal{A})^{**} \rightarrow V^\perp$ and so there is an $\mathcal{A}$ -bimodule

isomorphism $\iota : (I - P)(\mathcal{A} \hat{\otimes} \mathcal{A})^{**} \rightarrow V^*$. For every $v \in V$ and $T \in V^*$ by definition of τ we have $\tau(T) = x + V^\perp$ for some $x \in (\mathcal{A} \hat{\otimes} \mathcal{A})^{**}$, thus

$$\langle \iota^{-1}(T), v \rangle = \langle (I - P)x, v \rangle = \langle x, v \rangle - \langle Px, v \rangle = \langle x, v \rangle = \langle x + V^\perp, v \rangle = \langle \tau(T), v \rangle,$$

that is, $\iota^{-1}(T)|_V = T$ for every $T \in V^*$. Now suppose that $(F_\alpha) \subseteq V^*$ is an approximate $(\mathcal{A} \hat{\otimes} \mathcal{A})_{ess}^*$ -virtual diagonal for $\mathcal{A}$ and set $M_\alpha = \iota^{-1}(F_\alpha) \in (\mathcal{A} \hat{\otimes} \mathcal{A})^{**}$ for every α . Then

$$(3.1) \quad a \cdot M_\alpha - M_\alpha \cdot a = \iota^{-1}(a \cdot F_\alpha - F_\alpha \cdot a) \rightarrow 0.$$

Moreover, for $a \in \mathcal{A}$ by Cohen's factorization theorem, there exist $x, b, y \in \mathcal{A}$ such that $a = ybx$. Since for every $\psi \in \mathcal{A}^*$ we have $bx \cdot \psi \cdot y \in \mathcal{A}_{ess}^*$ and since $\mathcal{A}_{ess}^* = (\Delta^*)^{-1}(V)$ [2, Lemma 7.7], $\Delta^*(bx \cdot \psi \cdot y) \in V$. But $M_\alpha|_V = \iota^{-1}(F_\alpha)|_V = F_\alpha$ for each α . Using (3.1)

$$(3.2) \quad \begin{aligned} \langle \Delta^{**}(M_\alpha) \cdot a, \psi \rangle - \langle \Delta^{**}(F_\alpha) \cdot b, x \cdot \psi \cdot y \rangle &= \langle \Delta^{**}(M_\alpha) \cdot a, \psi \rangle - \langle F_\alpha, \Delta^*(bx \cdot \psi \cdot y) \rangle \\ &= \langle \Delta^{**}(M_\alpha) \cdot a, \psi \rangle - \langle M_\alpha, \Delta^*(bx \cdot \psi \cdot y) \rangle \\ &= \langle \Delta^{**}(M_\alpha) \cdot a, \psi \rangle - \langle \Delta^{**}(M_\alpha), bx \cdot \psi \cdot y \rangle \\ &= \langle \Delta^{**}(M_\alpha) \cdot a, \psi \rangle - \langle y \cdot \Delta^{**}(M_\alpha) \cdot bx, \psi \rangle \rightarrow 0. \end{aligned}$$

Also since (F_α) is an approximate $(\mathcal{A} \hat{\otimes} \mathcal{A})_{ess}^*$ -virtual diagonal and since $\Delta^*(x \cdot \psi \cdot y) \in V$, we have

$$(3.3) \quad \langle \Delta^{**}(F_\alpha) \cdot b, x \cdot \psi \cdot y \rangle \rightarrow \langle x \cdot \psi \cdot y, b \rangle.$$

Hence (3.2) and (3.3) imply that

$$(3.4) \quad \begin{aligned} |\langle \Delta^{**}(M_\alpha) \cdot a, \psi \rangle - \langle \psi, a \rangle| &\leq |\langle \Delta^{**}(M_\alpha) \cdot a, \psi \rangle - \langle \Delta^{**}(F_\alpha) \cdot b, x \cdot \psi \cdot y \rangle| \\ &\quad + |\langle \Delta^{**}(F_\alpha) \cdot b, x \cdot \psi \cdot y \rangle - \langle x \cdot \psi \cdot y, b \rangle| \rightarrow 0. \end{aligned}$$

We conclude from (3.1) and (3.4) that

$$a \cdot M_\alpha - M_\alpha \cdot a \rightarrow 0, \quad \langle \Delta^{**}(M_\alpha) \cdot a, \psi \rangle \rightarrow \langle \psi, a \rangle,$$

which means that (M_α) is an approximate virtual diagonal for $\mathcal{A}$. $\square$

As a corollary of Lemma 2.2, if a Banach algebra $\mathcal{A}$ is approximately amenable, then $F(\mathcal{A})$ is approximately Connes-amenable. For the rest of the paper we consider the converse of this result in the special case, whenever $\mathcal{A} = \ell^1(G)$. Indeed, we show that for a discrete group G , if $\ell^1(G)$ has an approximate WAP-virtual diagonal, then it has an approximate virtual diagonal, although the existence of an approximate virtual diagonal is weaker than approximate amenability of $\ell^1(G)$.

We identify $L^1(G) \hat{\otimes} L^1(G)$ with $L^1(G \times G)$ as $L^1(G)$ -bimodules (see [1, Example VI. 14] for instance).

Let I be the closed ideal in $L^\infty(G \times G)_{ess}$ generated by $\Delta^*(C_0(G))$ as in [2, Definition 8.1], that is,

$$I = \overline{\text{lin}\{\Delta^*(f)g : f \in C_0(G), g \in L^\infty(G \times G)_{ess}\}}.$$

Remark 3.3. The space of uniformly continuous bounded functions on G is denoted by $\text{UCB}(G)$. Since $\Delta^*(\text{UCB}(G)) = \Delta^*(L^\infty(G)_{ess}) \hookrightarrow L^\infty(G \times G)_{ess}$, we obtain a map $\Delta_{\text{UCB}} : L^\infty(G \times G)_{ess}^* \rightarrow \text{UCB}(G)^*$ as the adjoint of the inclusion map. By [2, Lemma 8.2], $\Delta^*(C_0(G)) \subseteq I \subseteq \text{WAP}(L^\infty(G \times G))$, so we

obtain a map $\Delta_{C_0} = (\Delta^*|_{C_0(G)})^* : I^* \rightarrow C_0(G)^* = M(G)$. By [2, Proposition 8.5] there is a G_d -bimodule map $S : I^* \rightarrow L^\infty(G \times G)_{ess}^*$ such that for every $h \in \text{UCB}(G)$ and $\psi \in I^*$

$$\langle \Delta_{\text{UCB}}(S(\psi)), h \rangle = \langle \iota(\Delta_{C_0}(\psi)), h \rangle,$$

where $\iota : M(G) \rightarrow \text{UCB}(G)^*$ is the natural inclusion map defined by $\langle \iota(\mu), h \rangle = \int_G h \, d\mu$ for every $\mu \in M(G)$ and $h \in \text{UCB}(G)$.

Theorem 3.4. *Let G be a discrete group. Consider the following statements*

- (i) $\ell^1(G)$ has an approximate WAP-virtual diagonal;
- (ii) $\ell^1(G)$ has an approximate $\ell^\infty(G \times G)_{ess}$ -virtual diagonal;
- (iii) $\ell^1(G)$ has an approximate virtual diagonal.

Then the implications (i) $\Rightarrow$ (ii) $\Rightarrow$ (iii) hold.

Proof. (i) $\Rightarrow$ (ii) Let $(M_\alpha) \subseteq F(\ell^1(G \times G))$ be an approximate WAP-virtual diagonal for $\ell^1(G)$. Then for every $a \in \ell^1(G)$

$$a \cdot M_\alpha - M_\alpha \cdot a \rightarrow 0 \quad (a \in \ell^1(G)) \quad \text{and} \quad \Delta_{\text{WAP}}(M_\alpha) \rightarrow e,$$

where e denotes the identity element of $F(\ell^1(G))$. Since $\Delta^*(c_0(G)) \subseteq I \subseteq \text{WAP}(\ell^\infty(G \times G))$, each M_α can be restricted to a functional on I . Let $S : I^* \rightarrow \ell^\infty(G \times G)_{ess}^*$ be as mentioned in Remark 3.3. Set $N_\alpha = S(M_\alpha|_I)$. We show that $(N_\alpha)_\alpha$ is an approximate $\ell^\infty(G \times G)_{ess}$ -virtual diagonal. Since G is discrete and S is a continuous G -bimodule map, it is a continuous $\ell^1(G)$ -bimodule map, so

$$a \cdot N_\alpha - N_\alpha \cdot a = S(a \cdot M_\alpha - M_\alpha \cdot a) \rightarrow 0 \quad (a \in \ell^1(G)).$$

Since (M_α) is an approximate WAP-virtual diagonal $\Delta_{c_0}(M_\alpha|_I) \cdot a \rightarrow a$ in $\ell^1(G)$ and since the map $\iota : \ell^1(G) \rightarrow \text{UCB}(G)^*$ is w^* - w^* -continuous, for every $h \in \text{UCB}(G)$ we have

$$\begin{aligned} \langle h, a \rangle &= \lim_\alpha \langle \iota \Delta_{c_0}(M_\alpha|_I \cdot a), h \rangle \\ &= \lim_\alpha \langle \Delta_{\text{UCB}} S(M_\alpha|_I \cdot a), h \rangle \\ &= \lim_\alpha \langle \Delta_{\text{UCB}}(N_\alpha \cdot a), h \rangle. \end{aligned}$$

Hence $(N_\alpha)_\alpha$ is an approximate $\ell^\infty(G \times G)_{ess}$ -virtual diagonal for $\ell^1(G)$.

(ii) $\Rightarrow$ (iii) holds by Proposition 3.2. □

Note that, it is not clear for us that the map S in Remark 3.3 is always an $L^1(G)$ -bimodule map. If S is an $L^1(G)$ -bimodule map, then the previous theorem holds not only for discrete groups but also for each locally compact group G .

REFERENCES

- [1] F. F. Bonsall, J. Duncan, *Complete Normed Algebra*, Vol. 80, Springer-Verlag, New York, 1973.
- [2] Y. Choi, E. Samei and R. Stokke, *Extension of derivations, and Connes-amenability of the enveloping dual Banach algebra*, arxiv:math.FA/1307.6287v2, 2014.
- [3] G. H. Esslamzadeh, B. Shojaei and A. Mahmoodi, *Approximate Connes-amenability of dual Banach algebras*, Bull. Belg. Math. Soc. Simon stevin **19** (2012), 193-213.

- [4] F. Ghahramani and R. J. Loy , *Generalized notions of amenability*, J. Funct. Anal. **208** (2004), 229-260.
- [5] B. E. Johnson, *Cohomology in Banach algebras*, Mem. Amer. Math. Soc. **127** (1972).
- [6] A.T.-M Lau and R. J. Loy, *Weak amenability of Banach algebras on locally compact groups*, J. Funct. Anal. **145** (1997), 175-204.
- [7] V. Runde, *Amenability for dual Banach algebras*, Studia Math. **148** (2001), 47-66.
- [8] ———, *Dual Banach algebras: Connes-amenability, normal, virtual diagonals, and injectivity of the predual bimodule*, Math. Scand. **95** (2004), 124-144.
- [9] ———, *Lectures on Amenability*, Lecture Notes in Mathematics, Vol. 1774, Springer-Verlag, Berlin, 2002.

FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, AMIRKABIR UNIVERSITY OF TECHNOLOGY, 424 HAFEZ AVENUE,
TEHRAN 15914, IRAN

E-mail address: shirinkalam_a@aut.ac.ir

E-mail address: arpabbas@aut.ac.ir